

Valid Names Results

Nipaecoccus neogaeus Williams & Granara de Willink, 1992 (Pseudococcidae: *Nipaecoccus*)

Nomenclatural History

- ***Nipaecoccus neogaeus*** Williams & Granara de Willink 1992: 274. Type data: TRINIDAD: Waller Field, on *Clidemia hirta*. Holotype, female, Type depository: London: The Natural History Museum, England, UK . accepted valid name Illustr.

Common Names

Ecological Associates

Hosts:

Families: 3 | Genera: 3

- Fabaceae
 - *Inga* | [BenDov1994](#) [WilliaGr1992](#)
- Malvaceae
 - *Theobroma bicolor* | [BenDov1994](#) [WilliaGr1992](#)
 - *Theobroma cacao* | [BenDov1994](#) [KondoRaVe2008](#) [WilliaGr1992](#)
- Melastomataceae
 - *Clidemia hirta* | [BenDov1994](#) [WilliaGr1992](#)

Geographic Distribution

Countries: 4

- Brazil
 - Bahia | [BenDov1994](#) [WilliaGr1992](#)
- Colombia | [BenDov1994](#) [Kondo2001](#) [KondoRaVe2008](#) [WilliaGr1992](#)
- Mexico | [BenDov1994](#) [WilliaGr1992](#)
- Trinidad and Tobago
 - Trinidad | [BenDov1994](#) [WilliaGr1992](#)

Keys

- [WilliaGr1992](#): pp.240 (Adult (F)) [Central and South America]

Remarks

- **General Remarks:** Good description and illustration of the adult female given by Williams & Granara de Willink (1992).

Illustrations

Citations

- [BenDov1994](#): 253
- [Kondo2001](#): distribution, host, taxonomy, 37

- [KondoRaVe2008](#): distribution, host, 38
- [WilliaGr1992](#): description, distribution, host, illustration, taxonomy, 274-277
- [WilliaMa2012](#): distribution, host, 88

[ABOUT](#) | [KEYS](#) | [KEY ASPIDIOTINI](#) | [CONTACT](#) | [LOGIN](#) | [FLAT CAT](#) | [CLASSIFICATION](#)

